



The effects of rootstock on the flower components of Clementine Mandarin (*Citrus clementina*)

Behzad Babazadeh Darjazi*

Department of Horticulture, Roudehen Branch, Islamic Azad University, Roudehen, Iran

Abstract

Studies have shown the importance of oxygenated compounds in beverage and food products. Citrus rootstocks seem to have a profound influence on oxygenated compounds in plants. The goal of the present study was to investigate the effect of rootstocks on the oxygenated compounds in *Citrus clementina*. Flower oil components were extracted using an ultrasonic bath and eluted with n-pentane: diethyl ether (1:2). The oils were then analyzed using GC and GC-MS. Data were analyzed using one-way analysis of variance (ANOVA) and Duncan's multiple range tests. Forty-one, 41, and 39 flower components were identified in Sour orange, Swingle citromelo and Troyer citrange, respectively. These included aldehydes, alcohols, ketones, monoterpenes, and sesquiterpenes. The major flavor components identified included linalool and sabinene. Among the three rootstocks examined, Swingle citromelo showed the highest content of aldehydes. Since aldehyde content of citrus is one of the most important indicators of quality, it seems that Citrus rootstocks have a profound influence on this factor.

Key words: Citrus rootstock; flavor components; flower oil

Babazadeh Darjazi, B. 2017. 'The effects of rootstock on the flower components of Clementine Mandarin (*Citrus clementina*)'. *Iranian Journal of Plant Physiology* 7 (2), 1999-2005.

Introduction

Mandarin is one of the most economically important crops in Iran. In the period 2011- 2012, the total mandarin production of Iran was estimated at around 825000 tonnes (FAO, 2012). *Citrus clementina* which is also called Yafa, is the most popular mandarin in the world. It is also one of the most important *Citrus* species cultivated in Iran and despite its importance, the flower components of *Citrus clementina* are relatively under-researched.

Citrus oils are commercially used for flavoring foods, beverages, perfumes, cosmetics, and medicines. In addition, recent studies have identified antimicrobial properties for Citrus oil (Babazadeh, 2009).

The quality of an essential oil can be calculated from the quantity of aldehyde compounds present in the oil. The quantity of aldehyde compounds present in the oil, is variable and depends upon a number of factors including rootstock (Verzera et al., 2003) and seasonal variation (Attaway et al., 1967) among other factors.

*Corresponding author

E-mail address: babazadeh @riau.ac.ir

Received: December, 2015

Accepted: July, 2016

Table 1
Common and botanical names for Citrus taxa used as scion and rootstock

Common name	Botanical name	Parents	Category
Clementine(scion)	<i>C. clementina</i> cv. <i>Cadox</i>	Unknown	Mandarin
Sour orange (Rootstock)	<i>Citrus aurantium</i> L.	Mandarin × Pomelo	Sour orange
Swingle citrumelo (Rootstock)	<i>Swingle citrumelo</i>	<i>C. paradisi</i> var <i>dancan</i> × <i>P. trifoliata</i> (L.) Raf	Poncirus hybrids
Troyer citrange (Rootstock)	<i>Troyer citrange</i>	<i>C. sinensis</i> × <i>P. trifoliata</i> (L.) Raf	Poncirus hybrids

Aldehydes are important flavor compounds extensively used in food products (Hemming, 2011). The quality of honey can be calculated from the amount of oxygenated components present in it (Alissandraki et al., 2003). In addition, type of flowers may influence the quality of volatile flavor components present in the honey. The effect of oxygenated compounds in the attraction of the pollinators has been established. Therefore, the presence of oxygenated compounds can encourage the agricultural yield (Kite et al., 1991).

In this paper, the flower compounds isolated from Clementine mandarin are compared with the aim of determining whether the quantity of oxygenated compounds is influenced by the rootstock.

Material and Methods

Clementine rootstocks

In 2007, rootstocks were planted at 4×4 m with three replications at an orchards in Ramsar (Latitude 36° 54' N, longitude 50° 40' E; Caspian Sea climate; average rainfall and temperature 970 mm and 16.25° C per year, respectively; soil was classified as loam-clay and pH ranged from 6.9 to 7). Sour orange, Swingle citrumelo and Troyer citrange were used as rootstocks in this experiment (Table 1).

Preparation of flower sample

Flowers were collected from many parts of the same trees in May 2015, early in the morning (6 to 8 am) and only during dry weather. The selection method of all samples was on a random basis.

Flower extraction techniques

In order to obtain the volatile compounds from the flowers, 50 g of fresh flowers were placed in a 2000 ml spherical flask, along with 300 ml of n-pentane:diethyl ether (1:2). The flask was covered and then placed in an ultrasonic water bath for 20 min. Extraction experiments were performed with an ultrasound cleaning bath-Fisatom Scientific-FS14H (Frequency of 40 KHz, nominal power 90 W and 24 × 14 × 10 cm internal dimensions water bath). The temperature of the ultrasonic bath was held constant at 25° C. The extract was subsequently filtered through MgSO₄ monohydrate. The extract was finally concentrated under a gentle stream of nitrogen to 1 ml and placed in a vial. Vial was sealed and kept in the freezer at -4° C until the GC-MS analysis (Alissandraki et al., 2003).

GC and GC-MS

An Agilent 6890N gas chromatograph (USA) equipped with a DB-5 (30 m × 0.25 mm i.d ; film thickness = 0.25 μm) fused silica capillary column (J&W Scientific) and a flame ionization detector (FID) was used. The column temperature was programmed from 60° C (3min) to 250° C (20 min) at a rate of 3° C/min. The injector and detector temperatures were 260° C and helium was used as the carrier gas at a flow rate of 1.00 ml/min and a linear velocity of 22 cm/s. The linear retention indices (LRIs) were calculated for all volatile components using a homologous series of n-alkanes (C9-C22) under the same GC conditions. The weight percent of each peak was calculated based on the response factor to the FID. Gas chromatography-mass spectrometry was used to identify the volatile components. The analysis was carried out with a Varian Saturn 2000R. 3800 GC linked with a Varian Saturn 2000R MS.

The oven condition, injector and detector temperatures, and column (DB-5) were the same as those given above for the Agilent 6890 N GC. Helium was the carrier gas at a flow rate of 1.1 ml/min and a linear velocity of 38.7 cm/s.




Fig. 1. HRGC chromatogram of flower oil of Clementine mandarin on Sour orange

Injection volume was 1 μ l.

Table 2
Flower components of Clementine mandarin on three different rootstocks

Component	Sour orange	Swingle citrumelo	Troyer citrange	Ki	Component	Sour orange	Swingle citrumelo	Troyer citrange	Ki
1 α -thujene	*	*	*	925	22 β - elemene	*	*	*	1343
2 α -Pinene	*	*	*	933	23 Cis-jasmone	*	*	*	1399
3 Benzaldehyde	*	*	*	954	24 (Z)- β -caryophyllene	*	*	*	1417
4 Sabinene	*	*	*	974	25 (Z)- β -farnesene	*	*	*	1451
5 β -Pinene	*	*	*	978	26 α - humulene	*	*	*	1461
6 β -myrcene	*	*	*	989	27 E,E- α -farnesene	*	*	*	1505
7 α -phellandrene	*	*	*	1003	28 (E)-Nerolidol	*	*	*	1562
8 δ -3-carene	*	*	*	1018	29 Caryophyllene oxide	*	*	*	1581
9 p-cymene	*	*	*	1024	30 Hexadecane	*	*	*	1593
10 Limonene	*	*	*	1030	31 Tetradecanal	*	*	*	1610
11 (Z)- β -ocimene	*	*	*	1035	32 8-heptadecene	*	*	*	1672
12 (E)- β -ocimene	*	*	*	1052	33 Pentadecanal	*	*	*	1690
13 γ - terpinene	*	*	*	1057	34 Heptadecane	*	*	*	1692
14 Octanol	*	*	*	1065	35 β -sinensal	*	*	*	1699
15 (E)-sabinene hydrate	*	*	*	1068	36 E,E-cis-farnesol	*	*	*	1731
16 α - terpinolene	*	*	*	1088	37 α -sinensal	*	*	*	1755
17 Linalool	*	*	*	1100	38 Caffeine	*	*	*	1849
18 Phenyl ethyl alcohol	*	*	*	1110	39 Nonadecane	*	*	*	1892
19 Terpinen-4-ol	*	*	*	1179	40 Eicosane	*	*	*	1992
20 α -terpineol	*	*	*	1192	41 Heneicosane	*	*	*	2094
21 Indol	*	*	*	1296		41	41	39	

*There is in oil

Identification of components

Components were identified by comparison of their Kovats retention indices (RI), retention times (RT), and mass spectra with those of reference compounds (Adams, 2001).

Data analysis

SPSS 18 was used for analysis of the data obtained from the experiments. Analysis of variations was based on the measurements of 7 flower components. Comparisons were made using one-way analysis of variance (ANOVA) and Duncan's multiple range tests. Differences were considered to be significant at $P < 0.01$. The correlation between pairs of characters was evaluated using Pearson's correlation coefficient.

Table 3

Statistical analysis of variation in flower components of Clementine mandarin on three different rootstocks

Compounds	Sour orange		Swingle citrumelo		Troyer citrange		F value
	Mean	St.err	Mean	St.err	Mean	St.err	
a) Aldehyds							
1) Benzaldehyde	0.20	0.02	0.30	0.03	0.20	0.02	
2) Tetradecanal	0.05	0.006	0.06	0.01			
3) Pentadecanal	0.03	0.006	0.03	0.006			
4) β -sinensal	1.97	0.12	2.24	0.09	1.41	0.10	F**
5) α -sinensal	2.45	0.11	2.63	0.10	2.13	0.13	F**
total	4.70	0.26	5.26	0.23	3.74	0.25	
b) Alcohols							
1) Octanol	0.10	0.006	0.10	0.01	0.10	0.01	
2) Linalool	20.91	0.27	26.00	0.20	15.72	0.18	F**
3) Phenyl ethyl alcohol	0.24	0.03	0.19	0.01	0.19	0.01	
4) Terpinen-4-ol	0.55	0.05	0.48	0.04	0.34	0.04	
5) α -terpineol	1.32	0.10	1.67	0.05	1.08	0.08	
6) Indol	0.90	0.05	0.80	0.05	0.80	0.05	
7) (E)-Nerolidol	2.78	0.10	3.11	0.11	2.48	0.09	F**
8) E,E-cis-farnesol	0.91	0.05	0.82	0.04	1.04	0.05	
total	27.71	0.65	33.17	0.52	21.75	0.52	
d) Ketones							
1) Cis-jasmone	0.27	0.02	0.23	0.02	0.17	0.02	
Monoterpenes							
1) α -thujene	0.07	0.006	0.07	0.01	0.08	0.01	
2) α -pinene	0.92	0.05	0.81	0.05	1.03	0.08	
3) Sabinene	21.52	0.20	19.23	0.23	24.84	0.20	F**
4) β -Pinene	1.47	0.09	1.54	0.10	1.87	0.09	
5) β -myrcene	2.34	0.10	2.19	0.11	2.48	0.11	
6) α -phellandrene	0.14	0.02	0.15	0.01	0.15	0.01	
7) δ -3-carene	0.75	0.05	0.78	0.04	0.88	0.07	
8) p-cymene	1.45	0.05	1.57	0.07	1.66	0.11	
9) Limonene	9.85	0.10	9.91	0.10	11.00	0.11	F**
10) (Z)- β -ocimene	0.18	0.02	0.18	0.01	0.20	0.02	
11) (E)- β -ocimene	6.31	0.10	7.04	0.09	8.96	0.10	F**
12) γ -terpinene	0.70	0.05	0.60	0.06	0.60	0.05	
13) (E)-sabinene hydrate	0.41	0.02	0.58	0.04	0.54	0.04	
14) α -terpinolene	0.37	0.02	0.31	0.02	0.36	0.04	
total	46.48	0.88	44.96	0.96	54.65	1.06	
Sesquiterpenes							
1) β -elemene	0.52	0.02	0.55	0.05	0.41	0.03	
2) (Z)- β -caryophyllene	0.99	0.05	1.15	0.08	1.64	0.08	
3) (Z)- β -farnesene	0.37	0.03	0.96	0.04	0.91	0.07	
4) α -humulene	0.07	0.01	0.06	0.01	0.06	0.01	
5) E,E- α -farnesene	0.26	0.02	0.27	0.02	0.24	0.02	
6) Caryophyllene oxide	0.38	0.04	0.41	0.03	0.35	0.05	
total	2.59	0.17	3.40	0.24	3.61	0.26	
1) Hexadecane	0.21	0.02	0.18	0.01	0.26	0.02	
2) 8-heptadecene	2.79	0.09	2.92	0.10	3.22	0.13	
3) Heptadecane	0.98	0.08	0.85	0.05	1.13	0.08	
4) Caffeine	0.11	0.01	0.22	0.02	0.13	0.01	
5) Nonadecane	1.26	0.05	0.97	0.07	1.36	0.08	
6) Eicosane	0.48	0.04	0.44	0.02	0.33	0.02	
7) Heneicosane	1.43	0.07	1.20	0.04	1.62	0.08	
total	7.26	0.38	6.78	0.31	8.05	0.43	
Total oxygenated compounds	32.68	0.94	38.66	0.77	25.66	0.79	
Total	89.01	2.38	93.80	2.28	91.97	2.54	

Mean is average composition (%) in three different rootstocks used with three replicates. St. err = standard error. F value is accompanied by its significance, indicated by: NS = not significant, * = significant at P = 0.05, ** = significant at P = 0.01.

Results

Flower components of the Clementine mandarin

GC-MS analysis of the flavor components extracted from Clementine mandarin using the ultrasonic bath allowed identification of 41 volatile components (Table 2, Fig. I) including 14 oxygenated terpenes (5 aldehydes, 8 alcohols, 1

ketone), 20 non-oxygenated terpenes (14 monoterpenes, 6 sesquiterpenes), and 7 other components.

Aldehydes

Five aldehyde components identified in this analysis were benzaldehyde, tetradecanal, pentadecanal, β -sinensal, and α -sinensal (Table 3). In addition, they were quantified from 3.74% to 5.26%. The concentration of α -sinensal was higher in the study samples. Among three rootstocks examined, Swingle citrumelo showed the highest content of aldehydes. Since the aldehyde content of Citrus oil is considered as one of the most important indicators of quality, rootstock apparently has a profound influence on this factor (Table 3).

Alcohols

Eight alcoholic components identified in this analysis were octanol, linalool, Phenyl ethyl alcohol, terpinen-4-ol, α -terpineol, indol, (E)-nerolidol and E,E-cis-farnesol (Table 3). The total concentration of alcohols ranged from 21.75% to 33.17%. Linalool was identified as the major component in this study and was the most abundant. Among three rootstocks examined, Swingle citrumelo showed the highest alcohol content (Table 3).

Ketones

One component identified in this analysis was cis-jasmone. The total amount of ketones ranged from 0.17% to 0.27%. Among three rootstocks examined, Sour orange showed the highest ketone content (Table 3).

Monoterpene hydrocarbons

The total amount of monoterpene hydrocarbons ranged from 44.96 % to 54.65 %. Limonene was identified as the major component in this study and was the most abundant. Among three rootstocks examined, Troyer citrange showed the highest content of monoterpenes (Table 3).

Sesquiterpene hydrocarbons

The total amount of sesquiterpene hydrocarbons ranged from 2.59% to 3.61 %. (Z)- β -caryophyllene was identified as the major component in this study and was the most abundant. Among three rootstocks examined, Troyer citrange showed the highest content of sesquiterpenes (Table 3).

Results of statistical analyses

Differences were considered to be significant at $P < 0.01$. These differences on the 1% level occurred in β -sinensal, α -sinensal, linalool, (E)-nerolidol, sabinene, limonene and (E)- β -ocimene (Table 3).

Results of correlation

Simple correlations between 6 components are presented in a correlation matrix (Table 4). The highest positive values of correlation coefficient (r) were observed between α -sinensal and β -sinensal. The highest significant negative correlations were observed between sabinene and linalool (Table 4).

Discussion

Our observation that rootstocks had an effect on some of the components of Clementine oil was in accordance with previous findings

Table 4

Correlation matrix (numbers in this table correspond with main components mentioned in Table 3)

	β -sinensal	α -sinensal	Linalool	(E)-Nerolidol	Sabinene	Limonene
α -sinensal	0.98**					
Linalool	0.97**	0.92**				
(E)-Nerolidol	0.97**	0.97**	0.91**			
Sabinene	0.94**	-0.94**	-0.97**	-0.91**		
Limonene	-0.85**	-0.75*	-0.94**	-0.27*	0.89**	
(E)- β -ocimene	-0.78*	-0.70*	-0.87**	-0.26*	0.77*	0.97**

*=significant at 0.05, **=significant at 0.01

(Verzera et al., 2003). Compositions of the flower oils obtained by ultrasonic bath from three rootstocks were very similar. However, the relative concentration of compounds was different according to the type of rootstock.

Comparison of our data with those in the literature revealed some inconsistencies with previous studies (Miguel. et al., 2008). This could be attributed to rootstock and environmental factors that can influence the compositions. However, it should be kept in mind that the extraction methods also may influence the results. Fertilizers and irrigation affect the content of oil present in Citrus (Kesterson et al., 1974). As fertilization, irrigation, and other operations were controlled in this study, this variability cannot be due to these factors.

The discovery of geranyl pyrophosphate (GPP), as an intermediate between mevalonic acid and oxygenated compounds (Alcohols and aldehyds), led to a rapid description of the biosynthetic pathway of oxygenated compounds. The biosynthetic pathway of oxygenated compounds in higher plants is as below:

Mevalonic acid → Isopentenyl Pyrophosphate → 3.3-dimethylallylpyrophosphate → geranyl pyrophosphate → Alcohols and Aldehyds

This reaction pathway is catalyzed by isopentenyl pyrophosphate isomerase and geranyl pyrophosphate synthase, respectively (Hay and Waterman, 1995). The pronounced enhancement in the amount of oxygenated compounds, when Swingle citrumelo was used as the rootstock, showed that either the synthesis of geranyl pyrophosphate was enhanced or activities of both enzymes increased.

Cytokinins can influence the essential oil components (Stoeva and Iliev, 1997). It is generally accepted that Cytokinins in higher plants are synthesized mainly in the root system and transported via the transpiration stream in xylem to the shoot. In addition, cytokinin level in the xylem sap also can vary by rootstock and exhibit a source-sink relationship by making strong sink tissues for mineral elements and

other metabolites including amino acids (Gordon et al., 1984).

High positive correlations between pairs of terpenes suggest a genetic control (Scora et al., 1976) and such dependence between pairs of terpenes was due to derivation of one from another that was not known. Similarly, high negative correlations between pairs of terpenes indicated that one of the two compounds had been synthesized at the expense of the other or of its precursor. Non-significant negative and positive correlations can imply genetic and/or biosynthetic independence. However, without an extended insight into the biosynthetic pathway of each terpenoid compound, the true significance of these observed correlations is not clear.

Considering that acetate is necessary for the synthesis of terpenes, it can be assumed that there is a specialized function for this molecule and it may be better served by Swingle citrumelo.

Conclusion

In the present study it was found that the amount of flower compositions was significantly affected by rootstocks and there was a great variation in most of the measured characters among three rootstocks. The study also demonstrated that volatile compounds in flower can vary when different rootstocks are utilized. Among three rootstocks examined, Swingle citrumelo showed the highest content of oxygenated compounds while the lowest oxygenated compound contents were produced by Troyer citrange. Further research on the relationship between rootstocks and oxygenated compounds is necessary.

Acknowledgements

The author thanks Roudehen Branch, Islamic Azad University for the financial support of the present research. Also the author would like to express his gratitude to Z. Kadkhoda from Institute of Medicinal Plants located at Supa Blvd-Km 55 of Tehran – Qazvin (Iran) for her help in GC-MS and GC analysis.

References

- Adams, R.P.** 2001. 'Identification of Essential Oil Components by Gas Chromatography / Mass Spectrometry'. Illinois: Allured Publishing Corporation. Carol Stream.
- Alissandrakis, E., D. Daferera, P.A. Tarantilis, M. Polissiou and P.C. Harizanis.** 2003. 'Ultrasound assisted extraction of volatile compounds from citrus flowers and citrus honey'. *Food Chemistry*, 82: 575-582.
- Attaway, J.A., A.P. Pieringer and L.J. Barabas.** 1967. 'The origin of citrus flavor components. III. A study of the percentage variation in peel and leaf oil terpenes during one season'. *Phytochemistry*, 6:25-32.
- Babazadeh Darjazi, B.** 2009. The effects of rootstock on the volatile flavor components of Page mandarin juice and peel, Agri. Sci. (Hort.) Dissertation. Islamic Azad University. Science and Research Branch, Tehran, Iran.
- Food and Agriculture Organization.** 2012. Statistics faostat-agriculture, production, crops, from <http://www.faostat.org>.
- Gordon, A., J. Stevens and N. W. Melvin,** 1984. 'Fruit set and cytokinin-like activity in the xylem sap of sweet cherry (*Prunus sativum*) as affected by rootstock'. *Physiologia Plantarum*, 61: 464-468.
- Hay, R.K.M. and P. Waterman.** 1995. Volatile Oil Crops: Their Biology, Biochemistry and Production. Wiley.
- Hemming, D.** 2011. Plant sciences reviews. CAB Reviews, CABI UK.
- Kesterson, J.W., R.J. Braddock and C.J. Koo.** 1974. 'The effect of bud wood, rootstock, irrigation and fertilization on the yield of Florida lemon oil', Proceeding Florida State Horticulture Society, pp: 6-9.
- Kite, G., T. Reynolds and T. Prance,** 1991. 'Potential pollinator –attracting chemicals from Victoria (Nymphaeaceae)'. *Biochemical Systematics and Ecology*, 19(7): 535-539
- Miguel, M.G., S.D. Andlen, A.C. Figueiredo, J.G. Barroso, L.G. Pedro, A. Duarte and J. Faisca.** 2008. 'Essential oils of flowers of *Citrus sinensis* and *Citrus clementina* cultivated in Algarve Portugal'. *Acta Horticulture*, 773: 89-94.
- Scora, R.W., A. Esen and J. Kumamoto.** 1976. 'Distribution of essential oils in leaf tissue of an F2 population of citrus'. *Euphytica*, 25: 201-209.
- Stoeva, T. and L. Iliev.** 1997. 'Influence of some phenyl urea cytokinins on spearmint essential oil composition'. *Bulgarian Journal of Plant Physiology*, 23(3):66-71.
- Verzera, A., A. Trozzi, F. Gazea, G. Ciccirello and A. Cotroneo,** 2003. 'Effect of rootstock on the composition of bergamot (*Citrus bergamia* Risso et Poiteau) essential oil'. *Journal of Agricultural and Food Chemistry*, 51: 206-210.

