



The effects of various levels of water deficit stress on forage yield and physiological characteristics of four cultivars of clover (*Trifolium* spp) under low input condition

Dariush Nematollahi¹, Hamid Reza Eisvand^{1*}, AliMohammad Modares Sanavi², Naser Akbari¹ and Ahmad Ismaili¹

1. Department of Agronomy and Plant Breeding, Faculty of Agriculture, Lorestan University, POB: 6815144316, Khorram Abad, Iran

2. Department of Agronomy and Plant Breeding, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

Abstract

This paper reports a study on the role of antioxidant enzymes and physiological characteristics of clover cultivars in their resistance against water deficit stress in the field conditions. Four cultivars of clover (*Trifolium* spp) namely, *T. resupinatum* L., *T. incarnatum* L., *Trifolium pretense* L., and *T. alexandrinum* L. under three levels of irrigation including %35, %50, and %65 soil moisture discharge were studied in a split plot experiment with completely randomized blocks design with three repetitions. Analysis of variance showed a significant effect of irrigation level on dry forage weight, leaf relative water content, and proline, catalase, peroxidase, chlorophyll b, and carotenoid contents. The effects of cultivars on dry forage weight, leaf relative water content, and proline, chlorophyll a, chlorophyll b, total chlorophyll, and carotenoid contents of the plants were also significant ($P \leq 0.01$). Generally, with an increase in the percentage of soil moisture, the dry forage yield showed a significant decrease. Also, water deficit resulted in increased proline content and also increased activities of catalase, peroxidase, and superoxide dismutase antioxidants. On the other hand, no significant differences were observed between cultivars with respect to the level of antioxidant activities. In addition, the effects of soil moisture discharge on the activities of all antioxidants except for those of superoxide dismutase were significant. Moreover, increasing the percentage of moisture discharge led to the reduction in the relative water content of the leaves and chlorophyll and carotenoid contents of all four cultivars of clover under study. Finally, the study concludes with recommending *T. incarnatum* L. as the most suitable clover cultivar for the climatic condition of the study region and under low input management (low irrigation, minimum fertilizer application, no herbicide application, and no weeding).

Keywords: antioxidant enzymes, water deficit, forage, clover

Nematollahi, D., H. R. Eisvand, A.M. Modares Sanavi, N. Akbari and A. Ismaili. 2017. ' The effects of various levels of water deficit stress on forage yield and physiological characteristics of four cultivars of clover (*Trifolium* spp) under low input condition'. *Iranian Journal of Plant Physiology* 7(3), 2125-2134.

*Corresponding author

E-mail address: eisvand.hr@lu.ac.ir

Received: January, 2016

Accepted: May, 2017

Introduction

Clover is a perennial forage plant belonging to the Papilionaceae family which has applications as fresh forage, pasture, silos, and also for modifying soil components (Futsaether et al., 2009). Because of nitrogen stabilizing nodules on their rhizome, clovers play an important role in soil enrichment and fertility and their fast and dense growth coupled with the potential for multiple harvest makes it possible for them to override and destroy weeds. Therefore, these plants are considered as an important option for alternation with other crops (Vaseva et al., 2011).

Due to severe limitations in water resources in most regions of Iran, water deficit stress is introduced as the most important type of stress on crops. The majority of the provinces except Gilan and Mazandaran are classified as arid and semiarid lands with unreliable precipitation where average annual rainfall is less than evaporation. Iran generally belongs to the regions where the moisture needs of clover during its growth should be supplied through irrigation water. Since in many regions of the country the rainfall may not be sufficient and because the growth stages are generally simultaneous with hot and dry summer weather condition when the water needs of other crops are also high, it is very likely that irrigation periods are lengthened or 2-3 irrigation turns are postponed during the growth stage (Kouchaki et al., 1997).

When water is a limiting factor in production of a crop, suitable irrigation treatment is needed to achieve maximum yield for consumption of each unit of irrigation water volume. Low irrigation is one of the methods of improving Irrigation Water Use Efficiency (IWUE) (Bekele and Tilahun, 2007). The main aim of low irrigation is increasing IWUE in crops through reducing the water volume in irrigation or decreasing the irrigation turns. In recent years, the strategies for improving efficiency in agricultural practices have centered on improving the efficiency of using limited available irrigation water resources with the effect that low irrigation is becoming a quite possible option (Garg and Dadhich, 2014).

As a result of increase in non-reproducible energy costs and also their damaging biological

effects, promotion of low input agriculture has grown considerably by government and non-government organizations in recent decades. The Since 1990, the United States has incorporated advising more low input and organic agricultural practices in its national programs with a view on the demands for these products in the market (Greene, 2013). Management methods are considered as an influential factor in the eventual efficiency of the farms. For example, introduction of groundcover plants and legumes in alternation with the main crops in the farm improves their yield and increases energy and nitrogen consumption efficiency. It must be noted that the introduction of fundamental changes in the main energy portals of the farm is time consuming. Also, replacement of the non-organic agricultural input has its own challenges and limitations. Therefore, increasing the efficiency of energy through low and high input management is a priority for many organizations that support the environment (Clerk et al., 2016).

Considering the increasing need of the society for healthy food and because of the limitations in water resources in the country and the high dependency of forage plants of the legume family to water, the capability of various cultivars of clover under different irrigation conditions is an important research issue. Moreover, livestock products are highly dependent upon forage plants of the legume family since legumes are an important resource for proteins in feeds. Therefore, taking into account the limitations in water resources, cutting the subsidies with a consequence of increased costs of agricultural inputs, and the increasing demands for healthy products, the need for research in this area is felt more than in the past.

Materials and Methods

This study was conducted in the research farm of Tarbiat Modares University of Tehran in a farm with an area of approximately 1000m² located in 35°, 74' N and 51°, 16'E. The farm was prepared for crop and in order to control the irrigation and level of drought stress, a drip irrigation system was

implemented in the farm using special PVC and plastic pipes for drip irrigation and the relevant connection.

The experiment was carried out based on a split plot experiment with completely random blocks with three repetitions where the irrigation and clover cultivar treatments were considered for the main and periphery plots, respectively. The study treatments consisted of three levels of irrigation (irrigation after discharging %35, %50, and %65 available soil moisture) and four cultivars of clover including *T. resupinatum* L., *T. incarnatum* L., *Trifolium pretense* L., and *T. alexandrinum* L. Seeds were obtained from Karaj Seed and Plant Improvement Institute. Nitrogen was supplied as urea (NH₂)₂CO as a starter, 20 kg/ha for all plots. No other chemical or organic fertilizers were applied in the study. Time Domain Reflectometry method (TDR) was used for deciding on the irrigation time.

During sowing, the seeds of *T. resupinatum* L., *T. incarnatum* L., *Trifolium pretense* L., and *T. alexandrinum* L. cultivars were inoculated with *Rhizobium leguminosarum* (Bovary trifolii). Seeds were sown in late April 2015. Irrigation was done based on moisture discharge. Low irrigation started from 4-leaf stage to 6-leaf stage based on the soil moisture discharges under study and continued up to harvest stage. Each plot had four lines of m with a distance of 50 cm. The seeds sown per hectare were 20 kg for *T. resupinatum* L. and *Trifolium pretense* L., and 25 kg for *T. alexandrinum* L. and *T. incarnatum* L. Twenty kg/ha nitrogen was used from the fertilizer tank as the starter. Soil profile of the study area is shown in Table 1.

In order to determine the qualitative yield of the forage, the plants were cropped from the two central lines of the plots disregarding those grown by 0.5 m of both ends which covered a total area of 2m² at %20 flowering stage and the harvest was immediately weighed and the fresh forage yield was recorded in each plot per kilogram. A sample (1 kg) of the fresh harvest was then

randomly selected, transported to the laboratory analysis, and dried at 75° C for 24 h in order to determine the dried weight. To determine dry forage yield, the total yield of different harvests was considered as the criterion. Farm management was implemented based on the low agricultural input. No herbicide was used and the weeds were controlled manually. Also, the RWC was measured using the method of Ritchie et al. (1990) based on the following equation:

$$RWC = [(FW - DW) / (SW - DW)] \times 100$$

Where FM is fresh leaf weight immediately after sampling, DW is the dry leaf weight after oven treatment, and SW is saturated leaf weight after immersion in distilled water.

Activities of catalase, peroxidase, and superoxide dismutase antioxidant enzymes were measured based on the methods reported by Dhindsa et al. (1981), Chance and Maehly (1995), and Laspina et al. (2005), respectively. Also, the proline content of the plants under different treatments was obtained based on the method of Bates et al. (1973).

Chlorophyll and carotenoid assay

Lichtenthaler and Wellburn (1983) was used for chlorophyll and carotenoid assay. In this method, 0.25 g leaf tissue was homogenized in a porcelain mortar containing 5 mL acetone %80. The solution was then submitted to spectrophotometry and at 663, 646, and 470 nm to measure chlorophyll a, chlorophyll b, total chlorophyll, and carotenoid contents based on the following equations (Lichtenthaler and Wellburn, 1983):

$$Chl\ a = (12.25\ A_{663} - 2.79\ A_{646}),$$

$$Chl\ b = (21.21\ A_{646} - 5.1\ A_{663}),$$

$$Chl\ t = Chl\ a + Chl\ b, \text{ and}$$

$$Car = (1000\ A_{470} - 1.8\ Chl\ a - 85.02\ Chl\ b)/198.$$

Table 1
Some physicochemical properties of the soil in depth of 60 cm

| Ec (dS.m ⁻¹) | pH | Clay% | Silt% | N% | N% | N% | P (ppm) | K (ppm) |
|--------------------------|------|-------|-------|----|------|-------|---------|---------|
| 1.2 | 7.67 | 10 | 18 | 72 | 0.12 | 1.267 | 36 | 209 |

Statistical analysis of the data and testing their normality were carried out using Microsoft Excel. Graphs were prepared using SAS 9.1 and the means were compared with Duncan multiple range test at $P \leq 0.05$.

Results

Dry forage yield

Results of ANOVA showed significant differences under cultivar, irrigation, and also interaction of effects of cultivar-irrigation in yields of clover (Table 2). Maximum dry forage yield (4411.7 kg/ha) was observed in cultivar *T. incarnatum* L. after %35 soil moisture discharge while minimum dry forage yield (406.6 kg/ha) was recorded in cultivar *Trifolium pretense* L., after %65 soil moisture discharge. Increase in soil moisture discharge level reduced yields in all clover cultivars; however, the reduction was not significant between %50 and %65 moisture discharge levels in cultivar *T. alexandrinum* L. (Table 3).

Relative water content

The effects of cultivars and irrigation levels on RWT were significant at $P \leq 0.05$ and $P \leq 0.01$, respectively while the interaction of effects of cultivars and irrigation levels was not significant (Table 2). Maximum percentage of RWT. Maximum leaf RWC was recorded under %35 soil moisture discharge treatment in *T. incarnatum* L. Increase in the percentage of soil moisture discharge reduced leaf RWC in all clover cultivars under study; however, the reduction at %50 moisture discharge in *T. incarnatum* L. was not statistically significant compared to %65 moisture discharge. Also the minimum leaf RWC was observed in *Trifolium pretense* L. under %65 moisture discharge (Table 3).

Proline content

Variations in the content of the amino acid proline were not similar in different clover cultivars and under increased drought treatment. Increase in soil moisture discharge from %35 to %65 led to a significant increasing trend in the proline content of the plants under study. Maximum proline content was recorded in cultivar

T. incarnatum L. under %65 soil moisture discharge treatment while this was not significantly different from the proline content under %50 discharge. On the other hand, minimum proline content was measured in cultivar *Trifolium pretense* L. under %35 soil moisture discharge treatment which showed no significant difference with cultivar *T. resupinatum* L. under %35 (Table 3).

Antioxidant enzymes activities

ANOVA showed that the effect of irrigation treatment was statistically significant on the activities of catalase and peroxidase enzymes while it had no significant effect on the activities of superoxide dismutase. The effects of various cultivars of clover on the activities of three enzymes were not significant; however, the interaction of effects of cultivar and irrigation treatments were significant in the plants under study (Table 2).

T. resupinatum L. showed the highest catalase activities under %65 soil moisture discharge which was not significantly different from the treatment of %50 moisture discharge. Maximum peroxidase activity was also observed in *T. incarnatum* L. cultivar under %50 soil moisture discharge whereas the %65 moisture discharge in clover cultivar *T. incarnatum* L. had the highest activity of superoxide dismutase. Generally, increase in the soil moisture discharge led to increased antioxidant activities in all cultivars under study (Table 4).

Chlorophyll and carotenoid contents

Various levels of irrigation had significant effects on chlorophyll b and carotenoid contents of the cultivars under study while the effect of this treatment on chlorophyll a and total chlorophyll contents of the plants was not significant. Moreover, the effects of cultivars on chlorophyll a, chlorophyll b, total chlorophyll, and carotenoid contents were significant at $P \leq 0.01$. Also, the interaction of the effects of cultivar-irrigation was significant on chlorophyll a and b contents (Table 5). The maximum chlorophyll a content (25.46 mg per fresh leaf weight) was observed under %35 moisture discharge treatment in *T. resupinatum* L. showing no significant difference from *T. incarnatum* L. under the same level of irrigation.

Table 2

Analysis of variance for irrigation and cultivars effects on yield and physiological traits of *Trifolium species*

| S.O.V | Dry forage yield | RWC | Proline | CAT | POD | SOD |
|-------------|------------------|--------------------|---------|--------------------|--------------------|-------------------|
| Replication | 9685.7 | 49.4 | 72.3 | 506.3 | 11.8 | 2.3 |
| Irrigation | 222198.3** | 983.5** | 604.0** | 51152.6** | 250.2** | 7.1 ^{ns} |
| error a | 3295.0 | 34.7 | 20.2 | 14.9 | 10.6 | 38.8 |
| Species | 120113.3** | 155.1* | 115.4* | 16.1 ^{ns} | 53.1 ^{ns} | 9.1 ^{ns} |
| I × S | 30148.5** | 25.9 ^{ns} | 14.8* | 167.6* | 47.0* | 99.9** |
| error b | 2431.2 | 21.9 | 15.0 | 0.1 | 0.2 | 2,14 |
| CV(%) | 29.11 | 12.76 | 11.37 | 13.49 | 13.43 | 6.28 |

ns, **, and *, indicate non-significant, significant at 1%, and significant at 5%, respectively.

Table 3

Comparison of mean dry forage yield, RWC, and proline content in clover species at different irrigation levels

| Species | Irrigation regime | Dry forage yield (kg/ha) | RWC (%) | Proline (µg g ⁻¹ FW) |
|------------------------|-------------------|--------------------------|---------|---------------------------------|
| <i>T. pratense</i> | 35% | 2678.8c | 52.74c | 0.1260cd |
| | 50% | 1785.8e | 42.25f | 0.1292c |
| | 65% | 406.6h | 33.28i | 0.1296c |
| <i>T. resupinatum</i> | 35% | 4020.3b | 58.58b | 0.1231d |
| | 50% | 2140.5d | 41.18f | 0.1261cd |
| | 65% | 1455.9f | 34.81h | 0.1576a |
| <i>T. incarnatum</i> | 35% | 4411.7a | 60.50a | 0.1269cd |
| | 50% | 2152.7d | 48.41d | 0.1573a |
| | 65% | 741.8g | 47.74d | 0.1600a |
| <i>T. alexandrinum</i> | 35% | 2698.6c | 52.52c | 0.1283cd |
| | 50% | 1233.4fg | 44.40e | 0.1344b |
| | 65% | 777.0g | 37.58g | 0.1372b |

Means with the same letter in each column have no significant difference (P<0.05).

The lowest chlorophyll a content in the study was recorded in *Trifolium pratense* L. cultivar under %65 soil moisture discharge treatment. Increase in soil moisture discharge significantly decreased chlorophyll a content in all cultivars under study. Moreover, the chlorophyll a content in *T. resupinatum* L. and *T. incarnatum* L., cultivars was higher than in *Trifolium pratense* L. and *T. alexandrinum* L. cultivars under all irrigation treatments and this dominance was not disturbed despite reduction under drought stress (Table 6).

Clover cultivar *T. incarnatum* L. under %35 soil moisture discharge condition contained significantly higher chlorophyll b contents as

compared with the other cultivars under all irrigation treatments. Because of higher level of the chlorophyll a compared with chlorophyll b, total chlorophyll showed a trend similar to that of chlorophyll a. The highest total chlorophyll content (35.83 mg per fresh leaf weight) was recorded in *T. incarnatum* L. cultivar under %35 soil moisture discharge treatment which was significantly different from other treatments. On the other hand, the lowest total chlorophyll content (13.42 mg per fresh leaf weight) was recorded in *Trifolium pratense* L. cultivar under %65 soil moisture discharge treatment. Generally,

increased drought stress significantly decreased total chlorophyll content in all clover cultivars under the study.

Variations in leaf carotenoid contents were less than those of chlorophyll a, b, and total under drought stress so that with an increase in soil moisture discharge from %35 to %50 in *T. incarnatum* L. cultivar and from %50 to %65 in *Trifolium pretense* L. and *T. alexandrinum* L. cultivars, no significant decrease was observed in leaf carotenoid contents of the clovers under investigation (Table 6).

Discussion

Production of dry matter and the growth pattern are determined by the two factors of temperature and genotype (Smith, 1970). Many studies have confirmed the interaction of effects of abiotic, genetic, and agricultural factors on the variations in yields of clover cultivars (Pankiw et al., 1977; Taylor, 1985). According to the findings of the study, it could be concluded that forage yield under drought stress reduces due to many reasons including reduction in the growth period of the plants and the resulting reduction in the light they receive, reduction in photosynthesis due to chlorophyll damage, increase in antioxidant activities and consequently increase in the energy consumed by the plants to confront drought, and many other known and unknown factors. Water deficit induces different effects in cell metabolism and stimulates accumulation of chaperonins and dehydrins in cell tissue to protect it from stress conditions (Vaseva et al., 2011).

Relative water content reduced as a result of drought stress in the study and this is also reported in *Rosmarinus officinalis* and *Triticum aestivum* under drought stress conditions (Gratani and Varone, 2004; Foulkes et al., 2007). Reduction in leaf RWC can occur as a result of decrease in percentage of the soil moisture and consequently decrease in leaf water potential due to increased drought stress. The reduction in the water content of plant tissues under drought stress can limit plant growth and incur physiological and metabolic changes in it (Mursalova et al., 2015). Sensitive cultivars produce more proline in comparison with

resistant cultivars during drought stress and as a result, these differences in the proline content among various plants are because of their genetic differences and the different strategies they use against stress condition. Increased level of this amino acid under stress condition can be attributed to its probable role in osmotic regulation (Singh et al., 2000). In fact, the role of proline in resistance against drought is very complicated so that some researchers consider increased level of proline content under drought stress merely due to a damage incurred by the stress. In reality, accumulation of proline under drought stress is a general reaction that occurs as a result of proline synthesis in the tissue, inhibition of proline oxidation, and prevention of its participation in protein synthesis (Pedrol et al., 2000). In a study on the effects of drought stress on antioxidant enzyme activities in clover, Antoline et al. (2010) reported that drought stress made a significant change in the activities of SOD, APX, and GR enzymes. Increase in catalase activities is also reported in alfalfa under water deficit conditions (Wang et al., 2008). The species with higher catalase activities were generally more resistant under water deficit condition and this suggests the role of catalase in the plants' resistance against oxidative stress resulting from drought (Table 4). Higher activity of peroxidase enzyme in plants depends on the mechanism of its operation in cells and their locus of production. Peroxidase not only plays a role in plants' resistance against drought through accumulation of peroxide hydrogen, but also it has a role in plant growth, suberification, cell connections, lignification, and detoxification of the toxic compounds all of which playing an important role in increasing plant resistance against drought stress (Ahmad and Parasad, 2012). Increased activity of peroxidase in alfalfa can be attributed to its role in changing electrical conductance of roots for transferring water and nutrients (Li et al., 2010). Superoxide dismutase is a powerful antioxidant which destroys the superoxide radical and therefore, it is a primary defense layer against free oxygen radicals (Alscher et al., 2002). Research shows the increased activity of this enzyme in plant cells in response to various abiotic stresses such as drought (Badawi et al., 2003). There are also studies reporting the increased

Table 4
Comparison of mean antioxidant enzymes activity in clover species at different irrigation levels

| Species | Irrigation regime | CAT (EU mg ⁻¹ protein min ⁻¹) | POD (EU mg ⁻¹ protein min ⁻¹) | SOD (EU mg ⁻¹ protein min ⁻¹) |
|------------------------|-------------------|---|---|---|
| <i>T. pratense</i> | 35% | 30.37f | 16.45i | 6.96gh |
| | 50% | 32.55f | 20.62h | 7.34fg |
| | 65% | 53.52c | 35.83f | 16.91c |
| <i>T. resupinatum</i> | 35% | 35.43e | 44.14e | 8.45f |
| | 50% | 80.21a | 44.42e | 12.13d |
| | 65% | 82.49a | 47.83 | 24.34b |
| <i>T. incarnatum</i> | 35% | 44.77d | 35.07f | 10.39e |
| | 50% | 67.04b | 25.52g | 10.82de |
| | 65% | 68.38b | 52.26c | 26.36a |
| <i>T. alexandrinum</i> | 35% | 36.39e | 23.05gh | 5.44h |
| | 50% | 40.90de | 70.85a | 9.94e |
| | 65% | 44.22d | 62.30b | 11.99d |

Means with the same letter in each column show no significant difference (P<0.05).

Table 5
Analysis of variance for irrigation and cultivars effects on physiological traits of Trifolium species

| S.O.V | Chlorophyll a | Chlorophyll b | Total Chlorophyll | Carotenoid |
|-------------|---------------------|--------------------|---------------------|------------------------|
| Replication | 24.1 | 1.3 | 44.0 | 1361.0 |
| Irrigation | 41.7 ^{ns} | 74.1 ^{**} | 70.4 ^{ns} | 132291.1 ^{**} |
| error a | 83.4 | 7.9 | 124.7 | 163.9 |
| Species | 185.6 ^{**} | 61.9 ^{**} | 390.0 ^{**} | 39786.6 ^{**} |
| I × S | 16.3 [*] | 25.1 ^{**} | 112.7 ^{ns} | 295.7 ^{ns} |
| error b | 27.7 | 7.8 | 48.4 | 106.9 |
| CV (%) | 28.31 | 14.50 | 27.86 | 15.9 |

ns, **, and *, indicate non-significant, significant at %1, and significant %5, respectively.

Table 6
Comparison of mean chlorophyll and carotenoid content in clover species at different irrigation levels

| Species | Irrigation | Chlorophyll a mg/g FW | Chlprpphyll b mg/g FW | Total chlorophyll mg/g FW | Carotenoid mg/g FW |
|------------------------|------------|--------------------------|--------------------------|------------------------------|-----------------------|
| <i>T. pratense</i> | 35% | 17.54e | 6.19c | 23.73e | 5.84d |
| | 50% | 14.02f | 3.47f | 17.49g | 4.32f |
| | 65% | 10.75h | 2.67h | 13.42i | 4.30f |
| <i>T. resupinatum</i> | 35% | 25.46a | 5.18d | 30.64c | 8.03a |
| | 50% | 22.33b | 4.36e | 26.69d | 6.27c |
| | 65% | 20.40d | 4.16e | 24.56d | 5.01e |
| <i>T. incarnatum</i> | 35% | 24.90a | 10.93a | 35.83a | 8.11a |
| | 50% | 21.60c | 9.67b | 31.27b | 7.81a |
| | 65% | 20.27d | 4.40e | 24.67e | 6.33bc |
| <i>T. alexandrinum</i> | 35% | 18.14e | 6.24c | 24.38e | 6.67b |
| | 50% | 14.39f | 5.95cd | 20.34f | 6.36bc |
| | 65% | 13.22g | 3.33g | 16.55h | 6.38bc |

Means with the same letter in each column show no significant difference (P<0.05).

antioxidant activities of the plants under drought stress in sunflower and alfalfa (Habibi et al., 2004; Antolin et al., 2010). In many plants, antioxidant mechanism is activated to reduce the damaging effects of free radicals resulted from the stress condition. Therefore, with an increase in the level of stress, so does the antioxidant activities of the plants. This approach shows the mechanism of

plants' tolerance against stress. The more compounds effective in plants' antioxidant activities, the higher plants' capability to neutralize the harmful effects of free oxygen radicals in peroxidizing lipids, proteins, and even nucleic acids and eventually the more tolerance against stress.

Chlorophyll content has always been used as a very useful criterion for evaluation of the plants' physiologic conditions. Studies have found variations in total chlorophyll content of different species under drought stress (Jiang and Haung, 2001). For example, Iannucci et al. (2002) found that leaf chlorophyll content in one-year-old clovers increased under moderate drought condition and with increase in stress level it significantly decreased. Reduced chlorophyll content under drought stress can be due to the stimulation of senescence even though the reaction of different plants would be different (Tize and Zeiger, 1991). Photosynthesis is influenced by drought stress in two ways: closing the stomata which results in CO₂ deficit for chloroplast activities, and the direct effect of drought which is observed as chlorophyll damage. Reactive species of oxygen produced in plants during drought stress can cause damage in photosynthesis system and eventually decompose chlorophyll. Drought stress was reported to reduce carotenoid contents of wheat and clover (Akhkha et al., 2011; Lazaridou and Koutroubas, 2004) which confirms the findings of this study.

Conclusion

All biochemical characteristics in the study were influenced by irrigation, cultivar, and interaction of effects of cultivar-irrigation. Soil moisture discharge increased proline contents and antioxidant enzyme activities in different cultivars of *Trifolium* which was dependent on the cultivar and the level of drought stress. On the other hand, with increasing soil moisture discharge, chlorophyll and carotenoid contents reduced. The findings showed that antioxidant enzymes play an effective role in the plant resistance against drought stress in clover cultivars and therefore have the potential for being utilized as biochemical indexes in clover cultivars. Comparison of the means in the characteristics under study revealed that in different soil moisture discharge treatments, cultivar *T. incarnatum* L. had the highest dry forage yield under low and moderate moisture discharge conditions whereas under high soil moisture discharge condition, *T. resupinatum* L., had the highest dry forage yield. Based on the findings of this study, regarding the limitations in water

resources in the country and also consumption of less chemical inputs, determination of cultivars resistant to low water conditions is of great significance in conservation of the ecosystem.

References

- Ahmad, P. and M.N.V. Parasad.** 2012. 'Abiotic stress responses in plants, metabolism, productivity and sustainability (eds), Springer, New York.
- Akhkha, A., T. Boutraa, and Y. A. Alhejely.** 2011. 'The rates of photosynthesis, chlorophyll content, dark respiration, proline and abscisic acid (ABA) in wheat (*Triticum durum*) under water deficit conditions'. *International Journal of Agricultural Biology*, 13:215- 223.
- Alscher, R.G., N. Erturk. and L.S. Heath.** 2002. 'Role of superoxide dismutases (SOD) in controlling oxidative stress in plant'. *Journal of Experimental Botany*, 153: 1331-1341.
- Antoline M.C., I. Muro and M. Sanchez-Diaz.** 2010. 'Application of sewage sludge improves growth photosynthesis and antioxidant activities of nodulated alfalfa plants under drought conditions'. *Environmental Experimental Botany*, 32: 68-75.
- Badawi, G.H., Y. Yamauchi, E. Shimada, R. Sasaki., N. Kawano, K., Tanaka and K. Tanaka.** 2003. 'Enhanced tolerance to salt stress and water deficit by overexpressing superoxide dismutase in tobacco (*Nicotiana tabacum*) chloroplasts'. *Plant Science*, 166: 919-928.
- Bates, L.R. Waldren and I. Teare.** 1973. 'Rapid determination of free proline for water-stress studies'. *Plant and soil*, 39: 205-207.
- Bekele, S. and K. Tilahun.** 2007. 'Regulated deficit irrigation scheduling of onion in a semiarid region of Ethiopia'. *Agric. Water Management*, 89:148-152.
- Chance, B. and A. Maehly.** 1955. 'Assay of catalases and peroxidases'. *Methods in enzymology*, 2:764-775.
- Dhindsa, R. S., P. Plumb-Dhindsa and T. A. Thorpe.** 1981. 'Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase'. *Journal of Experimental Botany*, 32: 93-101.

- Foulkes M. J., R. Sylvester-Bradley., R. Weightman., and J. W. Snape.** 2007. Identifying physiological traits associated with improved drought resistance in winter wheat. *Field Crops Research*, 103,11–24.
- Futsaether C.M., A. V. Vollsnes, O. M. Kruse, E. Otterholt, K. Kvaal and A. B. Eriksen.** 2009. 'Effects of the nordic photoperiod on ozone sensitivity and repair in different clover species studied using infrared imaging'. *Ambio*, 38: 437–443.
- Garg, N.K. and S. M. Dadhich.** 2014. 'Integrated non-linear model for optimal cropping pattern and irrigation scheduling under deficit irrigation'. *Agric. Water Management*,140: 1–13.
- Gratani, L. and L. Varone.** 2004. 'Leaf key traits of *Erica arborea* L., *Erica multifolia* L. and *Rosmarinus officinalis* L. co-occurring in Mediterranean marquis. *Flora - Morphology, Distribution, Functional Ecology of Plants*,199(1): 58-69.
- Greene, C.** 2013. 'Growth patterns in the U.S. organic industry.' Amber Waves,October 24. USDA Economic Research Service. <http://www.ers.usda.gov/amberwaves/2013-october/growth-patterns-in-the-us-organic-industry.aspx.VJGdbdLF9p5>.
- Habibi, D., M.R. Ardakani, M. M. Boojari, D. E. Taleghani and A. Mahmoodi.** 2004. 'Antioxidative in sunflower subjected to drought stress'. *4th International Crop Science Congress*. 26 sep to 10 Oct. Australia.
- Iannucci, A., M. Russo, L. Arena, N. Difonzo and P. Martiniello.** 2002. 'Water deficit effects on osmotic adjustment and solute accumulation in leaves of annual clovers'. *European Journal of Agronomy*, 16: 111-122.
- Jiang, Y. and N. Huang.** 2001. 'Drought and heat stress injury to two cool-season turf grasses in relation to antioxidant metabolism and lipid peroxidation'. *Crop Science*, 41: 436-442.
- Kouchaki, AS., A.S., Nakh Forosh and Zarif C. ketabi.** 1997. 'Agriculture organic (translation)'. Ferdowsi University of Mashhad Press.Iran.
- Laspina, N., M. Groppa, M. Tomaro and M. Benavides.** 2005. 'Nitric oxide protects sunflower leaves against Cd-induced oxidative stress'. *Plant Science*, 169: 323-330.
- Lazaridou M and S. D. Koutroubas.** 2004. 'Drought effect on water use efficiency of Berseem clover at various growth stages'. New directions for a diverse planet: Proceedings of the 4th International Crop Science Congress Brisbane, Australia, 26 Sept–1 Oct 2004.
- Li, W.R.,S. Q. Zhang, S. Y.Ding and L. Shan.** 2010. 'Root morphological variation and water use in alfalfa under drought stress'. *Acta Ecologia Sinica*, 30(19): 5140-5150.
- Lichtenthaler, H. and A. R. Wellburn.** 1983. 'Determination of total carotenoids and chlorophyll a and chlorophyll b leaf extracts in different solvents'. *Biochemistry*, 603: 591-592.
- Mursalova J., Z. Akparov., J. Ojaghi., M. Eldarov., S. Belen., N. Gummadov. and A. Morgounov.** 2015. 'Evaluation of drought tolerance of winter bread wheat genotypes under drip irrigation and rain-fed conditions'. *Turkish Journal of Agriculture and Forestry*, 39: 1–8.
- Pankiw, P., S. G. Bonina and J. A. C. Lieveise.** 1977. 'Effect of row spacing and seeding rates on seed yield in red clover, alsike clover and birds foot trifoli'. *Canadian Journal of Plant Science*, 57: 413-418.
- Pedrol, N., P. Ramos and M.J. Riegosa.** 2000. 'Phenotypic plasticity and acclimation to water deficits in velvet grass: a long-term greenhouse experiment. Changes in leaf morphology, photosynthesis and stress-induced metabolites'. *Plant Physiology*, 157: 383-393.
- Ritchie, S. W. and H. T. Nguyen.** 1990. 'Leaf water content and gas exchange parameters of two wheat genotypes differing in drought resistance'. *Crop Science*, 30: 105-111.
- Singh DK, PWG, Sale, CK., Pallaghy and V. Singh.** 2000. 'Role of proline and leaf expansion rate in the recovery of stressed white clover leaves with increased phosphorus concentration'. *New Phytologist*, 146:261–269.
- Smith. D.** 1970. 'Influence of temperature on the yield chemical composition of five forage legume species'. *Agronomy Journal*, 62:520-525.
- Taylor. N. L.** 1985. 'Clover science and technology'. American Society. Agronomy, Medison, WI.

Tize, L. and **E. Zeiger.** 1991. 'Plant Physiology: Mineral Nutrition'. The Benjamin Cummings Publishing Co., Inc., Redwood City, CA.

Vaseva, I. ,Y. Akiscan, Y. D., Klimentina and U. Feller. 2011. 'Drought stress tolerance of red and white clover—comparative analysis of some chaperonins and dehydrins'. *Scientia Horticulturae*, 130:653–659.

Wang, Ch.Q., YF, Zhang and Y.B. Zhang. 2008. 'Scavenger enzyme activities in subcellular fractions of white clover (*Trifolium repens* L.) under PEG-induced water stress'. *Journal of Plant Growth Regulation*, 27:387–393.